

ON THE GEOMETRY OF BIFURCATION CURRENTS FOR QUADRATIC RATIONAL MAPS.

FRANÇOIS BERTELOOT AND THOMAS GAUTHIER

ABSTRACT. We describe the behaviour at infinity of the bifurcation current in the moduli space of quadratic rational maps. To this purpose, we extend it to some closed, positive $(1, 1)$ -current on a two-dimensional complex projective space and then compute the Lelong numbers and the self-intersection of the extended current.

1. Introduction.

For any holomorphic family $(f_\lambda)_{\lambda \in M}$ of degree d rational maps on \mathbb{P}^1 , the bifurcation locus is the subset of the parameter space M where the Julia sets J_λ of f_λ does not move continuously with λ . In their seminal paper [MSS], Mañé, Sad and Sullivan have shown that the bifurcation locus is nowhere dense in M and coincides with the closure of the set of parameters for which f_λ admits a non-persistent neutral cycle.

For the quadratic polynomial family $(z^2 + c)_{c \in \mathbb{C}}$, the bifurcation locus is the boundary of the Mandelbrot set and, in particular, it is bounded. The situation is much more complicated for quadratic rational maps. Their moduli space \mathcal{M}_2 can be identified with the complement in a complex projective space \mathbb{P}^2 of a line at infinity \mathbb{L}_∞ around which the behaviour of the bifurcation locus is far from being completely understood.

The first results in this direction are due to J. Milnor [M] who studied the curves

$$\text{Per}_n(w) := \{[f] \in \mathcal{M}_2 \text{ s.t. } f \text{ has a } n\text{-cycle of multiplier } w\}$$

in the projective space \mathbb{P}^2 (see Proposition 2.3). Sharper results have then been obtained by A. Epstein [E] and yields a precise description of the intersections of $\text{Per}_n(w)$ with \mathbb{L}_∞ (see Proposition 4.1). It should be stressed that the deformations of the Mandelbrot set, provided by intersecting the bifurcation locus with the *lines* $\text{Per}_1(t)$ for $|t| < 1$ (see Proposition 2.4), play a central role in these investigations. In particular, the work of C. Petersen [P] on the collapsing of limbs (see the beginning of the fourth section) is crucial in Epstein's proof.

Currents not only provide an appropriate framework for studying bifurcations from a measure-theoretic viewpoint, but are also very well suited to investigate the asymptotic distribution of the hypersurfaces $\text{Per}_n(w)$. Let us recall that L. DeMarco has proved that the bifurcation locus of any holomorphic family supports a closed, positive $(1, 1)$ -current [DeM1]. This current is denoted T_{bif} and called the *bifurcation current*. We refer the reader to the survey [Du] or the lecture notes [Be] for a report on recent results involving bifurcation currents and further references.

The link between the bifurcation current T_{bif} and the hypersurfaces $\text{Per}_n(w)$ relies on the fact that the Lyapunov exponent $L(\lambda)$ of f_λ (with respect to its maximal entropy measure) is a global potential for T_{bif} (see [DeM2] or [BB1]). G. Bassanelli and the first author have

indeed deduced from this property the convergence of the weighted integration currents on $\text{Per}_n(w)$

$$\lim_n \frac{1}{d^n} [\text{Per}_n(w)] = T_{\text{bif}}$$

for $|w| < 1$ or for any $w \in \mathbb{C}$ when the hyperbolic parameters are dense in M (see [BB3]).

The present paper is devoted to the study of the behaviour at infinity of both the bifurcation current and its self-intersection in the moduli space \mathcal{M}_2 of quadratic rational maps. Our motivation is twofold. We first consider the bifurcation current with the aim to obtain a precise counterpart of Milnor's and Epstein's result in the setting of currents. We then investigate its self-intersection with the hope to apply this to the problem of the equidistribution of the hypersurfaces $\text{Per}_n(e^{i\theta})$ for a fixed θ . As we shall see, the results about the self-intersection are somehow surprising and lead to interesting new questions. Let us now discuss the content of the paper more precisely.

We show that T_{bif} extends naturally to a closed, positive $(1, 1)$ -current \hat{T}_{bif} on \mathbb{P}^2 . We relate this current with the hypersurfaces $\text{Per}_n(w)$ and precise its support (see Theorem 3.1). More precisely, we show that its support intersects the line \mathbb{L}_∞ along a segment $\mathbb{L}_{\infty, \text{bif}}$, which contains the intersection points $\{\infty_{p/q}\}_{p/q \in \mathbb{Q}^*}$ of the curves $\text{Per}_n(w)$ and the line \mathbb{L}_∞ as a dense subset.

We then use Epstein's result to compute the Lelong numbers of \hat{T}_{bif} at any point of the line at infinity \mathbb{L}_∞ and get a precise description of the measure $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ (see Theorem 4.2). Namely, we prove that this measure is totally discrete and supported by the points $\infty_{p/q}$'s. The ideal point $\infty_{p/q}$ also appears to be the point at which the limbs $\mathcal{L}_{p/q}$ and $\mathcal{L}_{-p/q}$ of the Mandelbrot set collapse under the motion given by the intersection of the bifurcation locus and the lines $\text{Per}_1(t)$, when $t \rightarrow e^{\pm 2i\pi p/q}$. We prove that the dirac mass of the measure $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ at the point $\infty_{p/q}$ coincides with the Lelong number of \hat{T}_{bif} at $\infty_{p/q}$ and is precisely given by the mass of $\mathcal{L}_{p/q} \cup \mathcal{L}_{-p/q}$ for the bifurcation measure of the family $(z^2 + c)_{c \in \mathbb{C}}$.

We finally describe the support of the so-called bifurcation measure $T_{\text{bif}} \wedge T_{\text{bif}}$ and compare this measure with $\hat{T}_{\text{bif}} \wedge \hat{T}_{\text{bif}}$ (see Theorem 5.1 and Proposition 5.2). We show that, in contrast to the case of cubic polynomials, the support of the bifurcation measure is unbounded. Let us stress out that the intersection \hat{T}_{bif}^2 at infinity is more complicated than expected regarding the complete description of \hat{T}_{bif} at infinity provided by Theorem 4.2. This fact invalidates our hope to apply some generalization of the so-called comparison principle for plurisubharmonic functions with Lelong numbers to the above mentioned problem of equidistribution for $\text{Per}_n(e^{i\theta})$.

We believe that these phenomena are related to the discontinuity of the dynamics at infinity within Milnor's compactification of \mathcal{M}_2 . It would certainly be interesting to make this point precise and to investigate this in relation with the dynamically natural compactification constructed by DeMarco (see [DeM3]).

Notation. The complex plane will be denoted by \mathbb{C} and the Euclidean unit disc in \mathbb{C} will be denoted by \mathbb{D} . The k -dimensional complex projective space is denoted by \mathbb{P}^k . A ball of radius r centered at x in some metric space is denoted by $B(x, r)$.

2. Preliminaries.

2.1. Lelong numbers and currents in \mathbb{P}^2 .

Let T be a closed, positive, $(1, 1)$ -current in \mathbb{P}^2 . The *mass* of T on an open set $U \subset \mathbb{P}^2$ is given by

$$\|T\|_U := \int_U T \wedge \omega$$

where ω is the standard Kähler form on \mathbb{P}^2 . When V is an algebraic curve in \mathbb{P}^2 and $[V]$ is the integration current on V , then $\|[V]\|_{\mathbb{P}^2} = \deg(V)$.

For any $x \in \mathbb{P}^2$, the *Lelong number* of T at x is given by

$$\nu(T, x) := \lim_{r \rightarrow 0} \frac{1}{\pi r^2} \|T\|_{B(x, r)}.$$

These numbers somehow measure the singularities of T . If V is an algebraic curve in \mathbb{P}^2 , then $\nu([V], x)$ is the multiplicity of V at x . We will mainly use the two following properties of Lelong numbers (see e.g. [Dem] Proposition 5.12 page 160 and Corollary 7.9 page 169).

Theorem 2.1. (1) *If $T_n \rightarrow T$, then for any $x \in \mathbb{P}^2$,*

$$\nu(T, x) \geq \limsup_{n \rightarrow +\infty} \nu(T_n, x).$$

(2) *If T_1 and T_2 are such that $T_1 \wedge T_2$ is well-defined, then for any $x \in \mathbb{P}^2$,*

$$\nu(T_1 \wedge T_2, x) \geq \nu(T_1, x) \cdot \nu(T_2, x).$$

2.2. The moduli space \mathcal{M}_2 of quadratic rational maps.

The space Rat_2 of quadratic rational maps on \mathbb{P}^1 may be viewed as a Zariski-open subset of \mathbb{P}^5 upon which the group of Möbius transformations acts by conjugation. The *moduli space* \mathcal{M}_2 is, by definition, the quotient resulting from this action. Denote by α, β, γ the multipliers of the three fixed points of $f \in \text{Rat}_2$ and $\sigma_1 = \alpha + \beta + \gamma$, $\sigma_2 = \alpha\beta + \alpha\gamma + \beta\gamma$ and $\sigma_3 = \alpha\beta\gamma$ the symmetric functions of these multipliers. Milnor has proved that (σ_1, σ_2) is a good parametrization of \mathcal{M}_2 (see [M]).

Theorem 2.2 (Milnor). *The map $[f] \in \mathcal{M}_2 \mapsto (\sigma_1, \sigma_2) \in \mathbb{C}^2$ is a biholomorphism.*

To study the curves

$$\text{Per}_n(w) := \{[f] \in \mathcal{M}_2 \text{ s.t. } f \text{ has a } n\text{-cycle of multiplier } w\},$$

it is useful to compactify \mathcal{M}_2 . In this context, it turns out that the projective compactification

$$[f] \in \mathcal{M}_2 \hookrightarrow [\sigma_1 : \sigma_2 : 1] \in \mathbb{P}^2$$

is suitable. Denote by \mathbb{L}_∞ the line at infinity of \mathcal{M}_2 , i.e.

$$\mathbb{L}_\infty = \{[\sigma_1 : \sigma_2 : 0] \in \mathbb{P}^2 / (\sigma_1, \sigma_2) \in \mathbb{C}^2 \setminus \{0\}\}.$$

Milnor has described the behavior of $\text{Per}_n(w)$ at infinity as follows (see [M] Lemmas 3.4 and section 4).

Proposition 2.3 (Milnor). (1) *For any $w \in \mathbb{C}$ the curve $\text{Per}_1(w)$ is a line in \mathbb{P}^2 whose equation in \mathbb{C}^2 is $(w^2 + 1)\lambda_1 - w\lambda_2 - (w^3 + 2) = 0$ and which intersects the line at infinity \mathbb{L}_∞ at the point $[w : w^2 + 1 : 0]$.*

- (2) For any $n \geq 2$ and any $w \in \mathbb{C}$, the curve $\text{Per}_n(w)$ is an algebraic curve in \mathbb{P}^2 whose degree equals the number $d(n) \sim 2^{n-1}$ of hyperbolic components of period n of the Mandelbrot set. In addition, the intersection $\text{Per}_n(w) \cap \mathbb{L}_\infty$ is contained in the set $\{[1 : u + 1/u : 0] \in \mathbb{P}^2 / u^q = 1 \text{ with } q \leq n\}$.

For reasons which will appear to be clear later, we shall denote by $\mathbb{L}_{\infty, \text{bif}}$ the subset of \mathbb{L}_∞ defined by

$$\mathbb{L}_{\infty, \text{bif}} := \{[1 : e^{i\theta} + 1/e^{i\theta} : 0] \in \mathbb{P}^2 / \theta \in [0, 2\pi]\}.$$

Golberg and Keen showed how the Mandelbrot set \mathbf{M} determines the connectedness locus for quadratic rational maps having an attracting fixed point (see [GK] Section 1 and [BB3] Theorem 5.4 for a potential theoretic approach).

Theorem 2.4 (Goldberg-Keen, Bassanelli-Berteloot). *There exists a holomorphic motion $\sigma : \mathbb{D} \times \text{Per}_1(0) \rightarrow \mathbb{C}^2$ such that $\mathbf{M}_t := \sigma_t(\mathbf{M}) \Subset \text{Per}_1(t) \cap \mathbb{C}^2$ is the connectedness locus of the line $\text{Per}_1(t)$. Moreover, if $|w| \leq 1$ and $n \geq 1$, then $\text{Per}_1(t) \cap \text{Per}_n(w) \subset \mathbf{M}_t$.*

2.3. Bifurcation currents in \mathcal{M}_2 .

Every rational map f of degree $d \geq 2$ on the Riemann sphere admits a unique maximal entropy measure μ_f . The Lyapunov exponent of f with respect to this measure is defined by

$$L(f) = \int_{\mathbb{P}^1} \log |f'| \mu_f.$$

It turns out that the induced function $L : \text{Rat}_2 \rightarrow \mathbb{R}$ is *p.s.h* and continuous on Rat_2 . We still denote by L the function induced on \mathcal{M}_2 . The *bifurcation current* T_{bif} is a closed positive $(1, 1)$ -current on \mathcal{M}_2 which may be defined by

$$T_{\text{bif}} = dd^c L.$$

As shown by DeMarco [DeM2], the support of T_{bif} coincides with the bifurcation locus of the family in the classical sense of Mañe-Sad-Sullivan (see also [BB1], Theorem 5.2). Using properties of the Lyapunov exponent, Bassanelli and the first author proved that the curves $\text{Per}_n(0)$ equidistribute towards the bifurcation current (see [BB3]).

Theorem 2.5 (Bassanelli-Berteloot). *The sequence $2^{-n}[\text{Per}_n(0)]$ converges to T_{bif} in the sense of currents in \mathcal{M}_2 . Moreover, $2^{-n}[\text{Per}_n(0)]|_{\text{Per}_1(0)}$ converges weakly to $\Delta(L|_{\text{Per}_1(0)}) = \frac{1}{2}\mu_{\mathbf{M}}$, where $\mu_{\mathbf{M}}$ is the harmonic measure of \mathbf{M} .*

As the function L is continuous, one can define the *bifurcation measure* μ_{bif} of the moduli space \mathcal{M}_2 as the Monge-Ampère mass of the function L , i.e.

$$\mu_{\text{bif}} := (dd^c L)^2 := dd^c (Ldd^c L).$$

This measure has been introduced by Bassanelli and the first author [BB1]. Buff and Epstein also studied it in [BE]. Recall that a rational map f is said to be strictly postcritically finite if each critical point of f is preperiodic but not periodic. We denote by \mathcal{SPCF} the set of classes of quadratic strictly postcritically finite rational maps. We also denote by \mathcal{S} the set of *Shishikura* maps, i.e. $\mathcal{S} := \{[f_0] \in \mathcal{M}_2 / f_0 \text{ has 2 distinct neutral cycles}\}$. Combining the work of Bassanelli and the first author with that of Buff and Epstein, we have the following.

Theorem 2.6 (Bassanelli-Berteloot, Buff-Epstein).

$$\text{supp}(\mu_{\text{bif}}) = \overline{\mathcal{SPCF}} = \overline{\mathcal{S}}.$$

These results are still valid in moduli spaces of degree d rational maps for any $d \geq 2$. In such a generality, this requires the result of [BG] that flexible Lattès examples belong to $\text{supp}(\mu_{\text{bif}})$.

3. Extension of the bifurcation current to Milnor's compactification.

As the current T_{bif} is a weak limit of weighted integration currents on curves which are actually defined on \mathbb{P}^2 , one may expect to naturally extend it to \mathbb{P}^2 . We will show how this is possible and prove some basic properties of the extended current. More precisely, we establish the following result which may be considered as a measure-theoretic counterpart of Milnor's description of the bifurcation locus in \mathcal{M}_2 .

Theorem 3.1. *There exists a closed positive $(1, 1)$ -current \hat{T}_{bif} on \mathbb{P}^2 whose mass equals $1/2$ and such that:*

- (1) $\hat{T}_{\text{bif}}|_{\mathbb{C}^2} = T_{\text{bif}}$,
- (2) $2^{-n}[\text{Per}_n(0)]$ converges to \hat{T}_{bif} in the sense of currents on \mathbb{P}^2 ,
- (3) $\text{supp}(\hat{T}_{\text{bif}}) = \text{supp}(T_{\text{bif}}) \cup \mathbb{L}_{\infty, \text{bif}}$.

We shall use the two following lemmas which are of independent interest. The first one is essentially due to Milnor (see Theorem 4.2 in [M] or Theorem 2.3 in [BB3]). The second one will be proved at the end of the section.

Lemma 3.2. $\|[\text{Per}_n(0)]\|_{\mathbb{P}^2} \sim 2^{n-1}$.

Lemma 3.3. $\overline{\bigcup_{n \geq 1} \text{Per}_n(0)} \cap \mathbb{L}_{\infty} = \mathbb{L}_{\infty, \text{bif}}$.

Proof of Theorem 3.1. We first justify the existence of \hat{T}_{bif} . According to the Skoda-El Mir Theorem (see [Dem] Theorem 2.3 page 139), the trivial extension of T_{bif} through the line at infinity \mathbb{L}_{∞} is a closed positive $(1, 1)$ -current on \mathbb{P}^2 if T_{bif} has locally bounded mass near \mathbb{L}_{∞} . The Lemma 3.2, combined with the fact that $2^{-n}[\text{Per}_n(0)]$ converges to T_{bif} on \mathbb{C}^2 (see Theorem 2.5), immediately yields $\|T_{\text{bif}}\|_{\mathbb{C}^2} \leq 1/2$. The extension \hat{T}_{bif} of T_{bif} therefore exists and $\|\hat{T}_{\text{bif}}\|_{\mathbb{P}^2} \leq 1/2$.

Let us now prove that $2^{-n}[\text{Per}_n(0)]$ converges to \hat{T}_{bif} on \mathbb{P}^2 and $\|\hat{T}_{\text{bif}}\|_{\mathbb{P}^2} = 1/2$. By Lemma 3.2, $(2^{-n}[\text{Per}_n(0)])_{n \geq 1}$ is a sequence of closed positive $(1, 1)$ -currents with uniformly bounded mass on \mathbb{P}^2 . According to the compactness properties of such families of currents, it suffices to show that \hat{T}_{bif} is the only limit value of $2^{-n}[\text{Per}_n(0)]$.

Assume that $2^{-n_k}[\text{Per}_{n_k}(0)]$ converges to T on \mathbb{P}^2 . By Siu's Theorem (see [Dem] Theorem 8.16 page 181), one has $T = S + \alpha[\mathbb{L}_{\infty}]$, where S has no mass on \mathbb{L}_{∞} . But, by Lemma 3.3, the line \mathbb{L}_{∞} is not contained in $\text{supp}(T)$. Thus $\alpha = 0$ and T has no mass on \mathbb{L}_{∞} . Since $T|_{\mathbb{C}^2} = \lim_k 2^{-n_k}[\text{Per}_{n_k}(0)] = T_{\text{bif}}$, this shows that T is actually the trivial extension of T_{bif} through \mathbb{L}_{∞} and therefore, according to the first part of the proof, equals \hat{T}_{bif} . Now, Lemma 3.2 immediately yields $\|\hat{T}_{\text{bif}}\|_{\mathbb{P}^2} = 1/2$.

It remains to prove that $\text{supp}(\hat{T}_{\text{bif}}) = \text{supp}(T_{\text{bif}}) \cup \mathbb{L}_{\infty, \text{bif}}$. The inclusion $\text{supp}(\hat{T}_{\text{bif}}) \subset \text{supp}(T_{\text{bif}}) \cup \mathbb{L}_{\infty, \text{bif}}$ follows immediately from items (1) and (2) and Lemma 3.3. The opposite inclusion is obtained from the fact that $\mathbb{L}_{\infty, \text{bif}} = \bigcup_{0 \leq \theta < 2\pi} (\text{Per}_n(e^{i\theta}) \cap \mathbb{L}_{\infty})$ and item (1), since $\text{Per}_n(e^{i\theta}) \cap \mathbb{C}^2 \subset \text{supp}(T_{\text{bif}})$. \square

Proof of Lemma 3.3. Let $\sigma : \text{Per}_1(0) \times \mathbb{D} \longrightarrow \bigcup_{|u|<1} \text{Per}_1(u)$ be the holomorphic motion given by Theorem 2.4. Let us set $\mathbf{M}_u := \sigma_u(\mathbf{M})$. As \mathbf{M} is compact and σ is continuous one has

$$\bigcup_{|u| \leq r < 1} \mathbf{M}_u \Subset \mathbb{C}^2 \text{ for all } 0 < r < 1.$$

Let us also recall that $\text{Per}_n(0) \cap \text{Per}_1(u) \subset \mathbf{M}_u$ for $n \geq 2$ and $u \in \mathbb{D}$.

We now proceed by contradiction and assume that there exists

$$\zeta \in (\overline{\bigcup_n \text{Per}_n(0)} \cap \mathbb{L}_\infty) \setminus \mathbb{L}_{\infty, \text{bif}}.$$

By Proposition 2.3, $\zeta \in \text{Per}_1(u_0)$ for some $u_0 \in \mathbb{D}$. Let us pick $\lambda_k \in \text{Per}_{n_k}(0)$ such that $\lambda_k \rightarrow \zeta$. Then there exists $u_k \in \mathbb{D}$ such that $u_k \rightarrow u_0$ and $\lambda_k \in \text{Per}_1(u_k)$, which is impossible since

$$\lambda_k \in \text{Per}_{n_k}(0) \cap \text{Per}_1(u_k) \subset \mathbf{M}_{u_k} \subset \bigcup_{|u| < r} \mathbf{M}_u$$

for some $|u_0| < r < 1$. □

4. Lelong numbers of the bifurcation current at infinity.

The aim of the present section is to compute the Lelong numbers of \hat{T}_{bif} at any point. This is related to previous works of Petersen ([P]) and Epstein ([E]) which we briefly describe.

Let \heartsuit be the main cardioid of the Mandelbrot set and $\mathcal{L}_{p/q}$ the p/q -limb of \mathbf{M} (see [Br] page 84). Denote by $d_{p/q}(n)$ the number of n -hyperbolic component of $\mathcal{L}_{p/q}$ and set

$$D_{p/q}(n) = \begin{cases} d_{p/q}(n) & \text{if } p/q = 1/2, \\ 2d_{p/q}(n) & \text{otherwise.} \end{cases}$$

Let σ be the holomorphic motion of $\text{Per}_1(0)$ given by Theorem 2.4 and

$$\infty_{p/q} := [1 : 2 \cos(2\pi p/q) : 0]$$

if $1 \leq p \leq q/2$ and $p \wedge q = 1$. Petersen has proved that the limb $\sigma_u(\mathcal{L}_{p/q})$ of \mathbf{M}_u disappears when u tends non-tangentially to $e^{-2i\pi p/q}$. Using this result, Epstein has precisely described the intersection $\text{Per}_n(w) \cap \mathbb{L}_\infty$. Namely, he proved the following:

Proposition 4.1 (Epstein). *For any $w \in \mathbb{C}$ and any $n \geq 2$,*

$$[\text{Per}_n(w)] \wedge [\mathbb{L}_\infty] = \sum_{\substack{1 \leq p \leq q/2 \leq n/2 \\ p \wedge q = 1}} \nu([\text{Per}_n(w)], \infty_{p/q}) \delta_{\infty_{p/q}} = \sum_{\substack{1 \leq p \leq q/2 \leq n/2 \\ p \wedge q = 1}} D_{p/q}(n) \delta_{\infty_{p/q}}.$$

From the above Proposition and by using the previous section, we deduce the following.

Theorem 4.2. (1) *The Lelong numbers of \hat{T}_{bif} are given by*

$$\nu(\hat{T}_{\text{bif}}, a) = \begin{cases} 1/6 & \text{if } a = \infty_{1/2}, \\ 1/(2^q - 1) & \text{if } a = \infty_{p/q} \text{ and } q \geq 3 \\ 0 & \text{if } a \in \mathbb{P}^2 \setminus \{\infty_{p/q} / p \wedge q = 1, 1 \leq p \leq q/2\}. \end{cases}$$

(2) *The measure $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ is discrete and given by*

$$\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty] = \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1}} \nu(\hat{T}_{\text{bif}}, \infty_{p/q}) \delta_{\infty_{p/q}}.$$

The proof requires the two following lemmas. The first one is a consequence of Theorem 3.1 and the second one relies on a simple computation. They will be proved at the end of the section.

Lemma 4.3. *The measure $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ is a well-defined positive measure on \mathbb{P}^2 of mass $1/2$.*

Lemma 4.4. *Let $\mu := \frac{1}{6}\delta_{\infty_{1/2}} + \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1, q \geq 3}} \frac{1}{2^q - 1} \delta_{\infty_{p/q}}$, then μ has mass $1/2$.*

Proof of Theorem 4.2. First observe that $\nu(\hat{T}_{\text{bif}}, a) = 0$ when $a \notin \mathbb{L}_{\infty, \text{bif}}$, since then \hat{T}_{bif} has a continuous potential in a neighborhood of a . Let us now pick $1 \leq p \leq q/2$ such that $p \wedge q = 1$. By item (3) of Theorem 3.1 and Theorem 2.1, we have

$$\nu(\hat{T}_{\text{bif}}, \infty_{p/q}) \geq \limsup_{n \rightarrow \infty} \nu(2^{-n}[\text{Per}_n(0)], \infty_{p/q}).$$

Proposition 4.1 states that $\nu([\text{Per}_n(0)], \infty_{p/q}) = D_{p/q}(n)$ is the number of n -hyperbolic components of the union $\mathcal{L}_{p/q} \cup \mathcal{L}_{-p/q}$ of limbs of the Mandelbrot set. Thus, by Theorem 2.5,

$$2^{-n} \nu([\text{Per}_n(0)], \infty_{p/q}) \longrightarrow \frac{1}{2} \mu_{\mathbf{M}}(\mathcal{L}_{p/q} \cup \mathcal{L}_{-p/q}).$$

On the other hand, Bullett and Sentenac have proved that $\mu_{\mathbf{M}}(\mathcal{L}_{p/q}) = \mu_{\mathbf{M}}(\mathcal{L}_{-p/q}) = \frac{1}{2^q - 1}$ (see [BS]). This gives $\nu(\hat{T}_{\text{bif}}, \infty_{p/q}) \geq \frac{1}{2^q - 1}$ if $p/q \neq 1/2$ and $\nu(\hat{T}_{\text{bif}}, \infty_{1/2}) \geq \frac{1}{2(2^2 - 1)} = \frac{1}{6}$.

Let $a \in \mathbb{L}_\infty$. By Theorem 2.1, we have $\nu(\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty], a) \geq \nu(\hat{T}_{\text{bif}}, a) \nu([\mathbb{L}_\infty], a)$. As \mathbb{L}_∞ is a line, we get $\nu(\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty], a) \geq \nu(\hat{T}_{\text{bif}}, a)$. For $a = \infty_{p/q}$, we thus have $\nu(\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty], \infty_{p/q}) \geq \frac{1}{2^q - 1}$ if $q \geq 3$ and $\nu(\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty], \infty_{1/2}) \geq \frac{1}{6}$, which we restate as

$$\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty] \geq \mu.$$

Since, according to Lemmas 4.3 and 4.4, the positive measures $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ and μ have the same mass, this yields $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty] = \mu$. We thus get item (2) and $\nu(\hat{T}_{\text{bif}}, \infty_{p/q}) = \frac{1}{2^q - 1}$ for $q \geq 3$ and $\nu(\hat{T}_{\text{bif}}, \infty_{1/2}) = \frac{1}{6}$.

From $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty] \geq \nu(\hat{T}_{\text{bif}}, a) \delta_a$ for any $a \in \mathbb{L}_\infty$ and $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty] = \mu$ we get $\nu(\hat{T}_{\text{bif}}, a) = 0$ for $a \neq \infty_{p/q}$. \square

Proof of Lemma 4.3. Let us first remark that, by Theorem 3.1, $\text{supp}(\hat{T}_{\text{bif}}) \cap \mathbb{L}_\infty = \mathbb{L}_{\infty, \text{bif}}$ and let us decompose \mathbb{P}^2 as the disjoint union of the line $\text{Per}_1(0)$ and on open set Ω which is biholomorphic to \mathbb{C}^2 . Let us stress that with these notations one has $\mathbb{L}_\infty \setminus \text{Per}_1(0) \cap \mathbb{L}_\infty \subset \Omega$. By Proposition 2.3, the set $\mathbb{L}_{\infty, \text{bif}}$ is compact in Ω . By definition, the current \hat{T}_{bif} has a potential u in Ω which is continuous in $\Omega \setminus \mathbb{L}_\infty$. By item (3) of Theorem 3.1, this potential is actually continuous in $\Omega \setminus \mathbb{L}_{\infty, \text{bif}}$. Let B_1 be a ball of Ω containing $\mathbb{L}_{\infty, \text{bif}}$ and $(B_i)_{i \geq 2}$ be a covering of $\Omega \setminus B_1$ by balls such that $\overline{B_i} \cap \mathbb{L}_{\infty, \text{bif}} = \emptyset$ for all $i \geq 2$.

As the set of points around which u is not locally bounded is contained in $\mathbb{L}_{\infty, \text{bif}}$ and as $\mathbb{L}_{\infty, \text{bif}} \cap \partial B_i = \emptyset$ and B_i is pseudoconvex for any $i \geq 1$, a result of Demailly asserts that $\hat{T}_{\text{bif}}|_{\Omega} \wedge [\mathbb{L}_\infty] = dd^c u \wedge [\mathbb{L}_\infty]$ is well-defined (see [Dem] Proposition 4.1 page 150). Since, by Proposition 2.3 and Theorem 4.2, \mathbb{L}_∞ and $\text{supp}(\hat{T}_{\text{bif}})$ don't intersect in a neighborhood of $\text{Per}_1(0)$, the measure $\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]$ is a well-defined positive measure on \mathbb{P}^2 .

Finally, Bézout's Theorem asserts that $\|\hat{T}_{\text{bif}} \wedge [\mathbb{L}_\infty]\|_{\mathbb{P}^2} = \|\hat{T}_{\text{bif}}\|_{\mathbb{P}^2} \cdot \|[\mathbb{L}_\infty]\|_{\mathbb{P}^2} = 1/2$. \square

Proof of Lemma 4.4. Denote by $\phi(n)$ the Euler function. As the sets $\{p \text{ s.t. } 1 \leq p \leq q/2, p \wedge q = 1\}$ and $\{q - p \text{ s.t. } 1 \leq p \leq q/2, p \wedge q = 1\}$ have same cardinality, we get

$$\begin{aligned} \mu(\mathbb{P}^2) &= \frac{1}{6} + \sum_{\substack{1 \leq p \leq q/2, \\ p \wedge q = 1, q \geq 3}} \frac{1}{2^q - 1} = \frac{1}{2(2^2 - 1)} + \frac{1}{2} \sum_{\substack{1 \leq p < q, \\ p \wedge q = 1, q \geq 3}} \frac{1}{2^q - 1} \\ &= \frac{1}{2} \sum_{q \geq 2} \left(\sum_{\substack{1 \leq p < q, \\ p \wedge q = 1}} \frac{1}{2^q - 1} \right) = \frac{1}{2} \sum_{q \geq 2} \frac{\phi(q)}{2^q - 1}. \end{aligned}$$

A classical result (see [HW] Theorem 309 page 258) asserts that the series $\sum_{n \geq 1} \phi(n) \frac{x^n}{1-x^n}$ locally uniformly converges on \mathbb{D} and that its sum is $\frac{x}{(1-x)^2}$. Therefore,

$$\mu(\mathbb{P}^2) = \frac{1}{2} \left(\sum_{q \geq 1} \frac{\phi(q)}{2^q - 1} - \phi(1) \right) = \frac{1}{2} \left(\sum_{q \geq 1} \frac{\phi(q) (\frac{1}{2})^q}{1 - (\frac{1}{2})^q} - \phi(1) \right) = \frac{1}{2}.$$

□

5. Behavior of the bifurcation measure near infinity.

One often compares the moduli space \mathcal{M}_2 of quadratic rational maps with the moduli space \mathcal{P}_3 of cubic polynomials. In this section, we enlight some important differences between these two spaces. We first show that the bifurcation measure is not compactly supported in \mathcal{M}_2 . Let us recall that the bifurcation measure is compactly supported and coincides with $\hat{\mu}_{\text{bif}}$ in \mathcal{P}_3 .

Theorem 5.1. *The cluster set of the support of μ_{bif} in \mathbb{P}^2 is precisely $\mathbb{L}_{\infty, \text{bif}}$.*

Proof. By Theorem 3.1, it suffices to show that $\mathbb{L}_{\infty, \text{bif}}$ is accumulated by points of $\text{supp}(\mu_{\text{bif}})$. Recall that $\text{Per}_1(e^{2i\pi\nu}) \cap \mathbb{L}_{\infty} = \{[1 : 2 \cos(2\pi\nu) : 0]\}$ for any $0 \leq \nu \leq 1$ (see Proposition 2.3). Let us fix $0 < \theta_0 < 1$. For $\theta > 0$ small enough, the lines $\text{Per}_1(e^{2i\pi\theta_0})$ and $\text{Per}_1(e^{2i\pi(\theta-\theta_0)})$ do not intersect on \mathbb{L}_{∞} and therefore,

$$\{\lambda(\theta)\} := \text{Per}_1(e^{2i\pi\theta_0}) \cap \text{Per}_1(e^{2i\pi(\theta-\theta_0)}) \subset \mathbb{C}^2.$$

Since $\lambda(\theta)$ has two distinct neutral fixed points, it belongs to $\text{supp}(\mu_{\text{bif}})$ (see Theorem 2.6). It remains to check that $\lim_{\theta \rightarrow 0} \lambda(\theta) = [1 : 2 \cos(2\pi\theta_0) : 0]$. By the holomorphic index formula (see [M]), the multiplier $\gamma(\theta)$ of the third fixed point of $\lambda(\theta)$ is given by

$$\gamma(\theta) = \frac{2 - (e^{2i\pi\theta_0} + e^{2i\pi(\theta-\theta_0)})}{1 - e^{2i\pi\theta}}.$$

As $\lim_{\theta \rightarrow 0} |\gamma(\theta)| = +\infty$, one has $\lim_{\theta \rightarrow 0} \|\lambda(\theta)\| = +\infty$. The conclusion follows, since $\lambda(\theta)$ stays on $\text{Per}_1(e^{2i\pi\theta_0})$ and $\text{Per}_1(e^{2i\pi\theta_0}) \cap \mathbb{L}_{\infty} = \{[1 : 2 \cos(2\pi\theta_0) : 0]\}$. □

We would like to extend μ_{bif} as a reasonable bifurcation measure on \mathbb{P}^2 . To this aim, we compare the trivial extension of μ_{bif} to \mathbb{P}^2 with $\hat{\mu}_{\text{bif}} := (\hat{T}_{\text{bif}})^2$. We prove the following.

Proposition 5.2. *There exists a positive measure μ_{∞} supported by $\mathbb{L}_{\infty, \text{bif}}$ which has a positive mass and such that*

$$\hat{\mu}_{\text{bif}} = \mu_{\text{bif}} + \frac{1}{36} \delta_{\infty_{1/2}} + \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1, q \geq 3}} \frac{1}{(2^q - 1)^2} \delta_{\infty_{p/q}} + \mu_{\infty}.$$

Let us stress that our previous results ensure the existence of μ_∞ as a positive measure. A recent result due to Kiwi and Rees concerning the number of (n, m) -hyperbolic components of \mathcal{M}_2 allows to see that μ_∞ has actually positive mass.

Proof. Arguing exactly as in the proof of Lemma 4.3, one sees that $\hat{\mu}_{\text{bif}} = \hat{T}_{\text{bif}} \wedge \hat{T}_{\text{bif}}$ is a well-defined positive measure on \mathbb{P}^2 . By definition, $\hat{\mu}_{\text{bif}}$ and μ_{bif} coincide on \mathbb{C}^2 . To prove the existence of μ_∞ , it thus only remains to justify that

$$\hat{\mu}_{\text{bif}} \geq \mu_{\text{bif}} + \frac{1}{36} \delta_{\infty_{1/2}} + \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1, q \geq 3}} \frac{1}{(2q-1)^2} \delta_{\infty_{p/q}}.$$

By Theorem 2.1 and Theorem 4.2, we have

$$\nu(\hat{\mu}_{\text{bif}}, \infty_{p/q}) \geq \nu(\hat{T}_{\text{bif}}, \infty_{p/q})^2 = \frac{1}{(2q-1)^2}, \text{ when } q \geq 3$$

and $\nu(\hat{\mu}_{\text{bif}}, \infty_{1/2}) \geq 1/36$. The existence of μ_∞ then follows, since

$$\hat{\mu}_{\text{bif}}|_{\mathbb{L}_\infty} \geq \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1}} \nu(\hat{\mu}_{\text{bif}}, \infty_{p/q}) \delta_{\infty_{p/q}}.$$

Let us now show that $\mu_\infty > 0$. One easily deduce from the convergence of $2^{-n}[\text{Per}_n(0)]$ towards T_{bif} in any family (see [BB3, Theorem 3.1]) that

$$\mu_{\text{bif}} = \lim_m \lim_n 2^{-(n+m)} [\text{Per}_n(0) \cap \text{Per}_m(0)]$$

on \mathbb{C}^2 . Thus $\mu_{\text{bif}}(\mathbb{C}^2) \leq \limsup_m \limsup_n 2^{-(n+m)} [\text{Per}_n(0) \cap \text{Per}_m(0)](\mathbb{C}^2)$. Kiwi and Rees [KR] proved recently that the number of (n, m) -hyperbolic components in \mathbb{C}^2 is at most

$$\left(\frac{5}{48} 2^n - \frac{1}{8} \sum_{q=2}^n \frac{\phi(q) \nu_q(n)}{2^q - 1} \right) 2^m + \mathcal{O}(2^m),$$

where $\nu_q(n) \sim 2^{n-1}/(2^q - 1)$ and $\nu_q(n) \geq 2^{n-1}/(2^q - 1) - 1/2$ (see [KR] Theorem 1.1). A standard transversality statement asserts that this number actually coincides with $[\text{Per}_n(0) \cap \text{Per}_m(0)](\mathbb{C}^2)$ (see [BB3] Theorem 2.8) when $n \neq m$. Thus

$$\mu_{\text{bif}}(\mathbb{C}^2) \leq \frac{5}{48} - \frac{1}{16} \sum_{q \geq 2} \frac{\phi(q)}{(2^q - 1)^2}.$$

Let us now proceed by contradiction, assuming that $\mu_\infty = 0$. Since \hat{T}_{bif} has mass $1/2$, Bézout's Theorem gives $\|\hat{\mu}_{\text{bif}}\|_{\mathbb{P}^2} = 1/4$. Therefore,

$$\frac{1}{4} = \|\mu_{\text{bif}}\| + \frac{1}{36} + \sum_{\substack{1 \leq p \leq q/2 \\ p \wedge q = 1, q \geq 3}} \frac{1}{(2^q - 1)^2} = \|\mu_{\text{bif}}\| + \frac{1}{2} \sum_{q \geq 2} \frac{\phi(q)}{(2^q - 1)^2} - \frac{1}{36},$$

which yields $\frac{25}{56} \leq \sum_{q \geq 2} \frac{\phi(q)}{(2^q - 1)^2}$. We then have

$$\frac{25}{56} \leq \sum_{q \geq 2} \frac{\phi(q)}{(2^q - 1)^2} \leq \frac{\phi(2)}{6} + \frac{\phi(3)}{28} + \frac{1}{8} \sum_{q \geq 4} \frac{\phi(q)}{2^q - 1}$$

which is impossible, since $\sum_{q \geq 1} \frac{\phi(q)}{2^q - 1} = 2$ (see proof of Lemma 3.2), $\phi(2) = 1$ and $\phi(3) = 2$. \square

References

- [Be] François Berteloot. Bifurcation currents in holomorphic families of rational maps. In *Pluripotential Theory*, volume 2075 of *Lecture Notes in Math.*, pages 1–93. Springer-Verlag, Berlin, 2013.
- [Br] Bodil Branner. The Mandelbrot set. In *Chaos and fractals (Providence, RI, 1988)*, volume 39 of *Proc. Sympos. Appl. Math.*, pages 75–105. Amer. Math. Soc., Providence, RI, 1989.
- [BB1] Giovanni Bassanelli and François Berteloot. Bifurcation currents in holomorphic dynamics on \mathbb{P}^k . *J. Reine Angew. Math.*, 608:201–235, 2007.
- [BB2] Giovanni Bassanelli and François Berteloot. Distribution of polynomials with cycles of given multiplier, 2009. preprint math.DS/0907.0011.
- [BB3] Giovanni Bassanelli and François Berteloot. Lyapunov exponents, bifurcation currents and laminations in bifurcation loci. *Math. Ann.*, 345(1):1–23, 2009.
- [BE] Xavier Buff and Adam L. Epstein. Bifurcation measure and postcritically finite rational maps. In *Complex dynamics : families and friends / edited by Dierk Schleicher*, pages 491–512. A K Peters, Ltd., Wellesley, Massachusetts, 2009.
- [BG] Xavier Buff and Thomas Gauthier. Perturbations of flexible Lattès maps. to appear in Bull. Soc. Math. France.
- [BS] Shaun Bullett and Pierrette Sentenac. Ordered orbits of the shift, square roots, and the devil’s staircase. *Math. Proc. Cambridge Philos. Soc.*, 115(3):451–481, 1994.
- [Dem] J.-P. Demailly. Complex analytic and differential geometry, 2011. free accessible book (<http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>).
- [DeM1] Laura DeMarco. Dynamics of rational maps: a current on the bifurcation locus. *Math. Res. Lett.*, 8(1-2):57–66, 2001.
- [DeM2] Laura DeMarco. Dynamics of rational maps: Lyapunov exponents, bifurcations, and capacity. *Math. Ann.*, 326(1):43–73, 2003.
- [DeM3] Laura DeMarco. The moduli space of quadratic rational maps. *J. Amer. Math. Soc.*, 20(2):321–355, 2007.
- [Du] Romain Dujardin. Bifurcation currents and equidistribution on parameter space, 2011. preprint math.DS/1111.3989.
- [E] Adam Lawrence Epstein. Bounded hyperbolic components of quadratic rational maps. *Ergodic Theory Dynam. Systems*, 20(3):727–748, 2000.
- [GK] Lisa R. Goldberg and Linda Keen. The mapping class group of a generic quadratic rational map and automorphisms of the 2-shift. *Invent. Math.*, 101(2):335–372, 1990.
- [HW] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. The Clarendon Press Oxford University Press, New York, fifth edition, 1979.
- [KR] Jan Kiwi and Mary Rees. Counting hyperbolic components, 2010. preprint arXiv:math.DS/1003.6104v1.
- [M] John Milnor. Geometry and dynamics of quadratic rational maps. *Experiment. Math.*, 2(1):37–83, 1993. With an appendix by the author and Lei Tan.
- [MSS] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. École Norm. Sup. (4)*, 16(2):193–217, 1983.
- [P] Carsten Lunde Petersen. No elliptic limits for quadratic maps. *Ergodic Theory Dynam. Systems*, 19(1):127–141, 1999.

François Berteloot, Université Paul Sabatier MIG. Institut de Mathématiques de Toulouse. 31062 Toulouse Cedex 9, France. *Email:* berteloo@picard.ups-tlse.fr

Thomas Gauthier, Université de Picardie Jules Verne, LAMFA UMR-CNRS 7352, 80039 Amiens Cedex 1, France, and, Stony Brook University, Institute for Mathematical Sciences, Stony Brook, NY 11794, USA. *Email:* thomas.gauthier@u-picardie.fr